



Apixaban in patients at risk of stroke undergoing atrial fibrillation ablation

Kirchhof, Paulus; Haeusler, Karl Georg; Blank, Benjamin F ; De Bono, Joseph; Callans, David; Elvan, Arif; Fetsch, Thomas; Van Gelder, Isabelle C; Gentlesk, Philip; Grimaldi, Massimo; Hansen, Jim; Hindricks, Gerhard; Al-Khalidi, Hussein R.; Massaro, Tyler ; Mont, Lluís; Nielsen, Jens Cosedis; Nolker, Georg; Piccini, Jonathan ; De Potter, Tom; Scherr, Daniel

DOI:

<https://academic.oup.com/eurheartj/advance-article/doi/10.1093/eurheartj/ehy176/4943979>

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kirchhof, P, Haeusler, KG, Blank, BF, De Bono, J, Callans, D, Elvan, A, Fetsch, T, Van Gelder, IC, Gentlesk, P, Grimaldi, M, Hansen, J, Hindricks, G, Al-Khalidi, HR, Massaro, T, Mont, L, Nielsen, JC, Nolker, G, Piccini, J, De Potter, T, Scherr, D, Schotten, U, Themistoclakis, S, Todd, DM, Vijgen, J & Di Biase, L 2018, 'Apixaban in patients at risk of stroke undergoing atrial fibrillation ablation', *European Heart Journal*.
<https://doi.org/https://academic.oup.com/eurheartj/advance-article/doi/10.1093/eurheartj/ehy176/4943979>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is a pre-copyedited, author-produced PDF of an article accepted for publication in *European Heart Journal* following peer review. The version of record [insert complete citation information here] is available online at: xxxxxxx [insert URL that the author will receive upon publication here].

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Apixaban in patients at risk of stroke undergoing atrial fibrillation ablation

Paulus Kirchhof MD¹⁻⁴, Karl Georg Haeusler MD^{4,5}, Benjamin Blank⁴, Joseph De Bono MD^{1,3}, David Callans MD⁶, Arif Elvan MD⁶, Thomas Fetsch MD⁸, Isabelle C. Van Gelder MD PhD⁹, Philip Gentlesk MD¹⁰, Massimo Grimaldi MD PhD¹¹, Jim Hansen MD¹², Gerhard Hindricks MD¹³, Hussein R. Al-Khalidi PhD¹⁴, Tyler Massaro PhD¹⁵, Lluís Mont MD¹⁶, Jens Cosedis Nielsen MD¹⁷, Georg Nölker MD¹⁸, Jonathan P. Piccini MD MHS^{15,19}, Tom De Potter MD²⁰, Daniel Scherr MD²¹, Ulrich Schotten MD PhD^{4,22}, Sakis Themistoclakis MD²³, Derick Todd MD²⁴, Johan Vijgen MD²⁵, Luigi Di Biase MD^{26,27}

1. University of Birmingham Institute of Cardiovascular Sciences
2. SWBH NHS Trust, Birmingham, UK
3. University Hospitals Birmingham, Birmingham, UK
4. Atrial Fibrillation NETwork association (AFNET), Münster, Germany
5. Center for Stroke Research Berlin & Department of Neurology, Charité – Universitätsmedizin Berlin, Germany
6. Hospital of the University of Pennsylvania, Philadelphia, USA
7. Isala Heart Center Zwolle, Zwolle, The Netherlands
8. The Clinical Research Institute, Munich, Germany
9. University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
10. Sentara Cardiovascular Research Institute, Norfolk, Virginia, USA
11. Ospedale Generale Regionale F. Miulli, Acquaviva delle Fonti, Italy
12. Gentofte Hospital, Hellerup, Denmark
13. Abteilung für Rhythmologie, Leipzig Heart Center, Leipzig, Germany
14. Department of Biostatistics & Bioinformatics, Duke University School of Medicine, USA
15. Duke Clinical Research Institute (DCRI), Durham, NC, USA
16. Hospital Clinic Barcelona, University of Barcelona, Barcelona, Spain
17. Department of Cardiology, Aarhus University Hospital, Denmark
18. Herz- und Diabeteszentrum NRW, Ruhr-Universität Bochum, Bad Oeynhausen, Germany
19. Duke University, Division of Cardiology Duke University Medical Center, Durham NC, USA
20. Cardiovascular Center, OLV Aalst, Belgium
21. Department of Cardiology, Medical University Graz, Austria
22. University Maastricht, Department of Physiology, Maastricht, Netherlands
23. Ospedale Dell'Angelo, Mestre, Italy
24. Liverpool Heart and Chest Hospital, Liverpool, UK
25. Jessa Ziekenhuis, Campus Virga Jesse, Hasselt, Belgium
26. Albert Einstein College of Medicine, at Montefiore Hospital, New York, USA;
27. Texas Cardiac Arrhythmia Institute at St. David's Medical Center, Austin, Texas, USA

Correspondence to

Professor Paulus Kirchhof
Institute of Cardiovascular Sciences, University of Birmingham and
SWBH and UHB NHS Trusts

IBR 136
Wolfson Drive
Birmingham, B15 2TT
United Kingdom
phone: +44 121 414 7042
e-mail: p.kirchhof@bham.ac.uk

word count (text incl. abstract and acknowledgments) : 3578 including disclosures, references: 1634

Abstract

Background. It is recommended to perform atrial fibrillation ablation with continuous anticoagulation. Continuous apixaban has not been tested.

Methods. We compared continuous apixaban (5 mg BD) to vitamin K antagonists (VKA, INR 2-3) in atrial fibrillation patients at risk of stroke a prospective, open, multi-center study with blinded outcome assessment. Primary outcome was a composite of death, stroke, or bleeding (BARC 2-5). A high-resolution brain magnetic resonance imaging (MRI) sub-study quantified acute brain lesions. Cognitive function was assessed by Montreal Cognitive Assessment (MoCA) at baseline and at end of follow-up.

Results. Overall, 674 patients (median age 64 years, 33% female, 42% non-paroxysmal atrial fibrillation, 49 sites) were randomized; 633 received study drug and underwent ablation; 335 undertook MRI (25 sites, 323 analyzable scans). The primary outcome was observed in 22/318 patients randomized to apixaban, and in 23/315 randomized to VKA (difference -0.38% [90% CI -4.0%, 3.3%], non-inferiority $p=0.0002$ at the pre-specified absolute margin of 0.075), including 2 (0.3%) deaths, 2 (0.3%) strokes, and 24 (3.8%) ISTH major bleeds. Acute small brain lesions were found in a similar number of patients in each arm (apixaban 44/162 (27.2%); VKA 40/161 (24.8%); $p=0.64$). Cognitive function increased at the end of follow-up (median 1 MoCA unit, $p=0.005$) without differences between study groups.

Conclusions. Continuous apixaban is safe and effective in patients undergoing atrial fibrillation ablation at risk of stroke with respect to bleeding, stroke, and cognitive function. Further research is needed to reduce ablation-related acute brain lesions. Funding: AFNET, DZHK, BMS/Pfizer; AXAFA-AFNET 5; NCT02227550.

Introduction

Catheter ablation is an effective¹⁻³ and increasingly used component of rhythm control therapy to improve symptoms in patients with atrial fibrillation.⁴⁻⁶ Atrial fibrillation ablation is associated with a risk of stroke and major bleeding.⁴⁻⁶ Continuous oral anticoagulation using vitamin K antagonists (VKA) such as warfarin can reduce the risk of embolic events to less than 1% when combined with peri-procedural heparin.⁷ Therefore, continuous oral anticoagulation is recommended in patients undergoing atrial fibrillation ablation.^{4, 6, 7} One randomized trial comparing rivaroxaban to warfarin in 218 patients found similar bleeding rates with rivaroxaban compared to warfarin: 21/114 (18.4%) patients with bleeding on rivaroxaban, 18/104 (17.3%) patients with bleeding on VKA, one patient with stroke.⁸ Another trial randomizing 635 atrial fibrillation ablation patients to dabigatran or VKA found 59/318 (18.6%) patients with bleeding on dabigatran, 54/317 (17%) patients with bleeding on VKA, and one patient with transient ischemic attack.⁹ Continuous apixaban has not been compared to VKA in atrial fibrillation ablation patients.

Atrial fibrillation ablation, unlike other ablation procedures, has been associated with declining cognitive function 90 days after the procedure, raising concerns about peri-procedural protection of the brain.^{10, 11} Furthermore, acute brain lesions without corresponding neurological symptoms are detected in ca. 25% of patients undergoing atrial fibrillation ablation by high resolution diffusion weighted brain magnetic resonance imaging (MRI), a sequence that detects acute cytotoxic brain edema.¹²⁻¹⁵ Cognitive function and acute brain lesions have not been evaluated in controlled clinical trials of patients undergoing atrial fibrillation ablation.

Objectives

Therefore, we conducted a randomized trial comparing continuous apixaban to continuous VKA therapy in patients undergoing atrial fibrillation ablation, including assessment of cognitive function in all patients and MRI-detected brain lesions in a sub-study.

Trial design

AXAFA – AFNET 5 (Anticoagulation using the direct factor *Xa* inhibitor apixaban during Atrial Fibrillation catheter Ablation: Comparison to vitamin K antagonist therapy) was an investigator-

initiated, prospective, parallel-group, randomized, open, blinded outcome assessment study comparing continuous apixaban therapy to vitamin K antagonist therapy. Details of the study design have been published.¹⁶ AXAFA – AFNET 5 was conducted in Europe and North America. The trial sponsor was AFNET, Münster, Germany (www.kompetenznetz-vorhofflimmern.de). AXAFA – AFNET 5 was designed by the steering committee in cooperation with AFNET and conducted in accordance with the declaration of Helsinki and the International Conference on Harmonization Good Clinical Practice Guidelines (ICH-GCP). The protocol was approved by ethical review boards at all institutions. The Clinical Research Institute (CRI, Munich, Germany) executed the study in cooperation with the steering committee and the sponsor. Data collection and entry was performed using the MARVIN® eCRF system.¹⁶⁻¹⁸ An independent steering committee and an independent data and safety monitoring board guided the trial. All serious adverse events were adjudicated by an independent endpoint review committee blind to study group and INR values. The Duke Clinical Research Institute served as the statistical core and performed the statistical analyses for the trial. The authors vouch for the accuracy and completeness of the data and for the fidelity of the trial to the protocol. This manuscript was written by the authors.

Study Population

AXAFA – AFNET 5 enrolled patients scheduled for a first atrial fibrillation ablation. Patients had at least one established stroke risk factor (age \geq 75 years, heart failure, hypertension, diabetes, or prior stroke). The full inclusion and exclusion criteria have been published (see **Table 1**).¹⁶

Treatment

At *baseline*, clinical parameters, stroke risk, heart rhythm, symptoms, quality of life (EQ-5D, SF-12¹, and Karnofsky performance status¹⁸), and cognitive function (Montreal Cognitive Assessment Test (MoCA))¹⁹ were assessed. Patients were randomized in a ratio of 1:1 to apixaban or vitamin K antagonist therapy (VKA). Randomization was stratified by study site and AF type (paroxysmal vs. persistent or long-standing persistent). The randomization scheme was generated via a computer program using permuted block of a random size.

Apixaban. Patients randomized to apixaban received 5 mg twice daily throughout the study period.

The apixaban dose was reduced to 2.5 mg twice daily if two or more of the following characteristics were present: age ≥ 80 years, body weight ≤ 60 kg, or serum creatinine level ≥ 1.5 mg/dL (133 $\mu\text{mol/L}$).^{16, 20} Apixaban was continued during the ablation procedure without interruption, including on the morning of ablation. Continuous anticoagulation in this group was defined as having taken all but one apixaban dose per week based on pill count.

VKA. Patients randomized to VKA were treated using the locally used VKA, e.g. warfarin, phenprocoumon, or acenocoumarol²¹, prescribed and dispensed following local routine. VKA therapy was monitored by international normalized ratio (INR) measurements; a minimum of three INR measurements was mandatory prior to ablation. The last INR prior to ablation needed to be 1.8 or higher. The time in the therapeutic range was calculated by the Rosendaal method.²² Continuous anticoagulation in this group was defined by therapeutic INR (INR ≥ 2) in all INR measurements 30 days prior to catheter ablation.

All patients underwent follow-up visits at the time of the ablation procedure and 3 months after ablation. At the *ablation visit*, continuous anticoagulation for at least 30 days prior to ablation was assessed and an ECG performed. Transesophageal echocardiography could be used following local practice. Interrupted anticoagulation required rescheduling of the ablation for 30 days unless (1) atrial thrombi were excluded by transesophageal echocardiogram and (2) effective anticoagulation was demonstrated prior to starting the ablation procedure by either taking at least two doses of apixaban (patients randomized to apixaban), or by an INR value ≥ 1.8 (patients randomized to VKA). A heparin bolus (100 IU/kg body weight) was required prior to or directly after transseptal puncture. The ablation procedure followed local practice and current guidelines.⁴⁻⁶ The protocol encouraged pulmonary vein isolation, the use of irrigated tip catheters, and flushing of all left atrial sheaths. Activated Clotting Time (ACT) was kept >300 s throughout the procedure. ACT measurements, details of the ablation technology used, delivered energy, procedure time, rhythm at beginning and end of procedure, and the need for cardioversion during the procedure were collected. An echocardiogram (transthoracic or intracardiac) was mandated directly after ablation to detect pericardial effusion.

At the 3 month visit, cognitive function and quality of life were reassessed, a 24 hour Holter ECG was performed, and study medication was returned. A *final phone call* to assess serious adverse events was performed 30 days after discontinuation of study drug.

MRI sub-study

Centres participating in the MRI sub-study (n=25) offered brain MRI to all eligible study patients. A brain MRI was performed within 48 hours after the ablation procedure. The MRI sequences were designed to detect all acute brain lesions, and to differentiate acute from chronic lesions. An imaging charta defined the MRI and adjudication workflow and brain MRI requirements (**supplementary table 2**). The following MRI-sequences were used: T2*-weighted imaging to screen for intracranial hemorrhage, diffusion-weighted imaging (DWI) and apparent diffusion coefficient (ADC) maps (post processed) to assess acute brain infarction, and Fluid-attenuated inversion recovery (FLAIR) to investigate the age of brain lesions.^{14, 15} DWI was conducted using a slice thickness of 2.5-3 mm (high resolution DWI) to enhance the sensitivity of MRI for small lesions.^{14, 15} Images failing the immediate quality check were repeated whenever feasible. All images were independently analyzed by two experienced neuro-radiologists blinded to treatment allocation.

Study Outcomes

The *primary outcome* measured from randomization was the composite of all-cause death, stroke, or major bleeding among modified intention-to-treat (mITT) population, defined as all randomized patients who received study drug and underwent catheter ablation. Safety was assessed in all randomized patients receiving study drug (safety population). Sensitivity analyses were performed in all randomized patients (ITT). Another sensitivity analysis compared events during the peri-ablation period defined from ablation to 7 days after the procedure.⁹ Major bleeding was defined according to the Bleeding Academic Research Consortium (BARC \geq 2).²³ All bleeding events were centrally adjudicated according to the BARC, ISTH, and TIMI classifications.^{23, 24}

Secondary Outcomes included time from randomization to ablation (ITT population), nights spent in hospital after ablation, activated clotting time (ACT) during ablation (summarized as median, 25th, 75th percentiles, and number of ACT measurements within the target range), all bleeding events, tamponade,

need for transfusion, and changes in quality-of-life and cognitive function compared to baseline. In the MRI sub-study, the prevalence and number of MRI-detected acute brain lesions were compared between groups.

Adverse events

All serious adverse events were collected, defined as adverse events that caused or prolonged hospitalization, caused disability or incapacity, were life-threatening, resulted in death or were important medical events. In addition, pregnancy, overdose, and cancer diagnosed after randomization were defined as serious adverse events. As AXAFA – AFNET 5 compared approved anticoagulants within their indications, non-serious adverse events were generally not reported, but those of special interest were defined and assessed. These comprised ablation-related complications including non-serious bleeding. The protocol encouraged brain imaging in patients who developed neurological abnormalities after the ablation procedure. All events from randomization to 3 months after index ablation procedure or to premature study termination were analyzed.

Statistical analysis

We estimated that a total of 650 patients (325 per group) were needed to detect a pre-specified margin of 7.5% (absolute difference) with 80% power using upper 1-sided 95% confidence interval (i.e., 2-sided 90% CI) with 3% attrition rate. The Farrington and Manning score test was used to compute sample size and power. The primary non-inferiority hypothesis was tested in the ablation population (mITT) using the method of Farrington and Manning score test with the pre-specified absolute margin of 0.075. In addition, a time-to-event analysis using Cox proportional hazards model with a relative margin of 1.44 was conducted. A multivariable Cox proportional hazards model controlling for the baseline risk factors of age, sex, weight, type of atrial fibrillation, and the CHADS2 factors was conducted. Changes in quality of life and cognitive function were assessed at 3 months compared to baseline using the EQ-5D and SF-12 questionnaires, Montreal Cognitive Assessment, and Karnofsky scale. Changes in quality of life were compared by Analysis of Covariance (ANCOVA) models including the treatment arms as an indicator variable and the baseline quality-of-life variables as covariates. To accrue sufficient events for a formal non-inferiority analysis, AXAFA – AFNET 5 was exclusively conducted in patients at risk of stroke

(**Table 1**) and counted bleeding events following the relatively broad BARC classification.²³ An independent data and safety monitoring board monitored the study for safety. The Haybittle–Peto boundary was used as stopping rule guidance.

Descriptive statistics for continuous and categorical variables were summarized as means (SDs) and medians (25th, 75th percentiles), and numbers (percentages), respectively. Comparisons between continuous variables were performed using the Wilcoxon rank-sum test or two-sample t-test depending on normality; comparisons between nominal variables were performed using the Pearson’s chi-square test or Fisher’s exact test, depending on expected cell sizes. All analyses were 2-sided and tested at the nominal 0.05 significance level. No adjustment was made for multiple testing. Statistical analyses were performed with SAS version 9.4 (SAS Institute Inc., Cary, NC).

Results

Trial Participants

AXAFA – AFNET 5 randomized 674 patients across 49 sites in 9 countries from February 2015 to April 2017. Overall, 633 patients took study drug and underwent atrial fibrillation ablation (mITT, ablation set, **Figure 1**). Demographic and clinical characteristics were well balanced between groups (**Table 1**). Transesophageal echocardiography was used in 549/633 (86.7%) patients. All or all but one apixaban doses per week were taken by 307/318 (97%) patients randomized to apixaban in the ablation set. The median time in therapeutic range in the 315 patients randomized to VKA in the ablation set was 84% (71, 97%). Time from randomization to ablation was not different between study groups (**Table 1**).

Primary outcome

Primary outcome events (BARC 2-5 bleeding, stroke, or death) were observed in 22/318 (6.9%) patients randomized to apixaban, and in 23/315 (7.3%) patients randomized to vitamin K antagonist therapy in the ablation set. Four events were classified as TIMI major bleeding, and 24 events are ISTH major bleeding (**Table 3**). Two patients died: one patient randomized to VKA, female, age 70, hypertensive, last blood pressure 156/76, last INR 2.6, underwent pacemaker implantation 8 days after ablation and experienced a massive intracerebral haemorrhage. Another patient randomized to apixaban, male, age 69, with paroxysmal atrial fibrillation, hypertension, heart failure, diabetes, and chronic obstructive

lung disease, was found dead in his bed 19 days after ablation without identifiable cause of death upon autopsy. Two patients randomized to apixaban had a stroke. Both had persistent AF and underwent transesophageal echocardiogram. One patient, male, age 63, hypertensive, ACT 236-398s, developed slurred speech with matching MRI lesion on the day of radiofrequency pulmonary vein isolation that fully resolved. Another patient, male, age 52, ACT 301-400s, hypertensive, developed weakness of the right arm with paresthesia of the right leg after cryo-balloon pulmonary vein isolation that persisted beyond hospital discharge. Tamponade occurred in 2 (apixaban) and 5 (VKA) patients and was managed by pericardial drainage and administration of protamine and vitamin K. One patient with tamponade in each study group received blood transfusions. Anticoagulants were continued in five patients with tamponade, and paused for 4 days in one patient randomized to apixaban, and for 8 days in one patient randomized to VKA. All patients were discharged from hospital and attended the 3 months follow up (n=6) or an end of study visit (n=1).

Apixaban was noninferior to VKA based on the non-inferiority margin of 7.5% (a difference of -0.38%, 90% confidence interval -4.0%, -3.3%, non-inferiority p=0.0002). Apixaban was also noninferior to VKA among all randomized patients as assessed by Cox proportional hazards model comparison between treatment groups using a relative non-inferiority margin of 1.44 (equivalent to 7.5% absolute; hazard ratio=0.88, 90% CI 0.55, 1.41, p=0.042, **Figure 2**). There was no statistical interaction between clinical stroke and bleeding risk factors and treatment groups (**Figure 3**).

Secondary outcome parameters

There was no difference in time to ablation or nights spent in hospital after the ablation between groups (**Table 4**). As expected, the last INR prior to ablation and activated clotting times achieved during ablation were lower in the patients randomized to apixaban (**Table 4**). Quality of life as assessed by the physical component of SF-12 (+2.5 (-2.1, 8.1) units) and Karnofsky scale (+10 (0,10)) improved during the study without differences between study groups (**Table 4**). At least mild cognitive dysfunction was found in 188/619 (30.4%) of the patients at baseline (pre-defined as MoCA<26, **Table 2**). At the end of follow-up, MoCA increased by a median of +1.0 (-1.0, 2.0) unit without differences between study groups, and 7.2% fewer patients had mild cognitive impairment (**Table 4**).

MRI sub-study

Acute brain MRI was performed in 335 patients across 25 centres. Clinical characteristics of the sub-study population were not different from the main study population, with the exception of a lower median weight in patients undergoing MRI (85.0 kg (74.5, 96.0)) compared to non-MRI patients (90.0 kg (80.0, 103.0)). Clinical characteristics were well balanced between MRI sub-study treatment groups. There were 323 analyzable MRIs. Acute brain MRI lesions (**Figure 4**) were found in 44/162 (27.2%) patients randomized to apixaban, and in 40/161 (24.8%) patients randomized to VKA ($p=0.635$), with very similar distribution of lesions between random groups (**Table 5**). Cognitive function at the end of follow-up was not different in patients with or without acute brain lesions (MoCA 27.1 ± 2.7 in 239 patients without MRI lesions, 27.1 ± 2.8 in 84 patients with MRI lesions, $p=0.91$).

Discussion

AXAFA – AFNET 5 demonstrated that continuous anticoagulation with apixaban is a safe and effective alternative to VKA in patients at risk of stroke undergoing atrial fibrillation ablation. AXAFA – AFNET 5 observed 4 TIMI major bleeding events in 633 patients (0.6%, **Table 3**) compared to 1 event in 248 patients in VENTURE-AF (0.4%)⁸. AXAFA – AFNET 5 observed 24 patients with ISTH major bleeding events (3.8%, **Table 3**) compared to 27 events in 635 patients in RE-CIRCUIT (4.3%)⁹. The numerical differences in ISTH major bleeding rates between AXAFA – AFNET 5 (apixaban 10 patients (3.1%); VKA 14 patients (4.4%); **Table 3**) and RE-CIRCUIT (dabigatran 5 patients (1.6%); VKA 22 patients (6.9%))⁹ could be due to chance variations in outcomes, differences in risk profile between the AXAFA – AFNET 5 and RE-CIRCUIT study populations, and due to the high time in therapeutic range in the VKA group in AXAFA – AFNET 5 (median TTR 84%). AXAFA – AFNET 5 included only patients with stroke risk factors, resulting in a mean CHA₂DS₂VASc score of 2.4 and a population that was 4-5 years older than in the published controlled trials in atrial fibrillation ablation.^{1-3, 8, 9} Despite the higher stroke risk, we observed few strokes: AXAFA – AFNET 5 found 2 strokes in 633 patients (0.3%), compared to 1 stroke in 248 patients in VENTURE-AF (0.4%)⁸, and 1 TIA in 635 patients in RE-CIRCUIT (0.2%)⁹. Equally, mortality was low (0.3%) and similar to VENTURE-AF (0.4%)⁸, RE-CIRCUIT (0%)⁹, and the EORP AF ablation registry (0.2%)²⁵.

AXAFA – AFNET 5 included 86 patients on acenocoumarol and 102 patients on phenprocoumon, 186 patients (29%) undergoing cryo-ablation³, and 84 patients undergoing atrial fibrillation ablation without transesophageal echocardiography without safety signals, providing some reassurance that these common patterns of clinical practice can be used on continuous apixaban or VKA therapy.^{4-6, 26}

The secondary outcomes observed in AXAFA – AFNET5 underpin the safety of continuous apixaban in atrial fibrillation ablation: time to ablation was not different between groups and quality of life and cognitive function improved equally in both study groups after ablation. High-resolution diffusion weighted brain MRI detected acute brain lesions at the expected rate (approximately 25%)^{12, 14, 15} without differences between study groups. Continuous anticoagulation does not fully prevent acute brain lesions, which can be caused by debris dislodging from ablation wounds, air emboli, or small thrombi.^{27, 28} Procedural improvements are desirable to reduce acute brain lesions during atrial fibrillation ablation.²⁹ Further analyses of the AXAFA – AFNET 5 data set may shed more light on risk factors for acute brain lesions in patients undergoing AF ablation on continuous anticoagulation. One prior study found reduced cognitive function 90 days after atrial fibrillation ablation on interrupted warfarin therapy compared to baseline.¹⁰ Reassuringly, cognitive function improved at the end of AXAFA – AFNET 5 without differences between study groups.

Limitations.

AXAFA – AFNET 5 was an open study, but with blinded outcome assessment. The non-inferiority margin was wide. The findings are consistent with prior studies with continuous dabigatran and rivaroxaban. While AXAFA – AFNET 5 was the first study comparing cognitive function after atrial fibrillation ablation in a controlled trial, the assessment was limited to global cognitive function. Differentiation between acute and chronic lesions was done by using an accepted combination of MRI sequences.^{14, 15}

Conclusions

Continuous apixaban therapy is a safe and effective alternative to VKA in patients at risk of stroke undergoing atrial fibrillation ablation with respect to stroke, major bleeding, cognitive function, and MRI-detected acute brain lesions.

Acknowledgments and disclosures

AXAFA - AFNET 5 is an investigator initiated trial. Sponsor of the trial is AFNET. AXAFA – AFNET 5 was partially funded by BMS/Pfizer, the DZHK (German Centre for Cardiovascular Research) and by the BMBF (German Ministry of Education and Research) to AFNET. Further support came from European Union (grant agreement No 633196 [CATCH ME] to BB, PK, LM, US, AFNET), British Heart Foundation (FS/13/43/30324 to PK), Leducq Foundation to PK, and the Netherlands Heart Foundation (RACE V) to US and IvG. The authors would like to thank Bianca-Maria Klein, Elisabeth Freund, and the study teams at CRI and AFNET for excellent project coordination, help with preparing the manuscript and ensuring complete accordance of the paper with the trial protocol and eCRF, and the members of the clinical trials statistics group at DCRI who provided the statistical analysis. Authors disclose financial links with Abbott/St Jude Medical (KGH, LM, JP, LDB), AFNET (TF, JP), Allergan (JP), ARCA Biopharma (JP), Bayer (KGH, JP, US), Biosense Webster (MG, JH, LDB), Biotronik (LM, LDB), Boehringer Ingelheim (KGH, ST) Boston Scientific (JdB, GH, LM, JP, LDB), Bristol-Myers Squibb/Pfizer (KGH, BB, ST), British Heart Foundation (PK), Clinical Research Institute Munich (JH), Daiichi Sankyo (KGH, ST), Edwards Lifesciences (KGH), EIP Pharma (KGH), European Union (PK), Getemed AG (KGH), Janssen Pharmaceuticals (JP), Johnson&Johnson (JH, LM), Medtronic (KGH, JDB, LM, JP, LDB), Novo Nordisk Foundation (JCN), Phillips (JP), Roche (US), Sanofi-Aventis (KGH, JP), Stereotaxis (LDB), Universita Svizzera Italiana (US). PK is inventor of two patents held by University of Birmingham.

References

1. Cosedis Nielsen J, Johannessen A, Raatikainen P, Hindricks G, Walfridsson H, Kongstad O, Pehrson S, Englund A, Hartikainen J, Mortensen LS, Hansen PS. Radiofrequency ablation as initial therapy in paroxysmal atrial fibrillation. *N Engl J Med* 2012;**367**(17):1587-95.
2. Verma A, Jiang CY, Betts TR, Chen J, Deisenhofer I, Mantovan R, Macle L, Morillo CA, Haverkamp W, Weerasooriya R, Albenque JP, Nardi S, Menardi E, Novak P, Sanders P. Approaches to catheter ablation for persistent atrial fibrillation. *N Engl J Med* 2015;**372**(19):1812-22.
3. Kuck KH, Brugada J, Furnkranz A, Metzner A, Ouyang F, Chun KR, Elvan A, Arentz T, Bestehorn K, Pocock SJ, Albenque JP, Tondo C, Fire, Investigators ICE. Cryoballoon or Radiofrequency Ablation for Paroxysmal Atrial Fibrillation. *N Engl J Med* 2016;**374**(23):2235-45.
4. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, Castella M, Diener HC, Heidbuchel H, Hendriks J, Hindricks G, Manolis AS, Oldgren J, Popescu BA, Schotten U, Van Putte B, Vardas P, Agewall S, Camm J, Baron Esquivias G, Budts W, Carerj S, Casselman F, Coca A, De Caterina R, Deftereos S, Dobrev D, Ferro JM, Filippatos G, Fitzsimons D, Gorennek B, Guenoun M, Hohnloser SH, Kolh P, Lip GY, Manolis A, McMurray J, Ponikowski P, Rosenhek R, Ruschitzka F, Savelieva I, Sharma S, Suwalski P, Tamargo JL, Taylor CJ, Van Gelder IC, Voors AA, Windecker S, Zamorano JL, Zeppenfeld K. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;**37**(38):2893-2962.
5. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC, Jr., Conti JB, Ellinor PT, Ezekowitz MD, Field ME, Murray KT, Sacco RL, Stevenson WG, Tchou PJ, Tracy CM, Yancy CW, Members AATF. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation* 2014;**130**(23):e199-267.
6. Calkins H, Hindricks G, Cappato R, Kim YH, Saad EB, Aguinaga L, Akar JG, Badhwar V, Brugada J, Camm J, Chen PS, Chen SA, Chung MK, Nielsen JC, Curtis AB, Davies DW, Day JD, d'Avila A, Natasja de Groot NMS, Di Biase L, Duytschaever M, Edgerton JR, Ellenbogen KA, Ellinor PT, Ernst S, Fenelon G, Gerstenfeld EP, Haines DE, Haissaguerre M, Helm RH, Hylek E, Jackman WM, Jalife J, Kalman JM, Kautzner J, Kottkamp H, Kuck KH, Kumagai K, Lee R, Lewalter T, Lindsay BD, Macle L, Mansour M, Marchlinski FE, Michaud GF, Nakagawa H, Natale A, Nattel S, Okumura K, Packer D, Pokushalov E, Reynolds MR, Sanders P, Scanavacca M, Schilling R, Tondo C, Tsao HM, Verma A, Wilber DJ, Yamane T. 2017 HRS/EHRA/ECAS/APHRS/SOLAECE Expert Consensus Statement on Catheter and Surgical Ablation of Atrial Fibrillation. *Heart Rhythm* 2017;**15**:e275-e444.
7. Di Biase L, Burkhardt JD, Santangeli P, Mohanty P, Sanchez JE, Horton R, Gallinhouse GJ, Themistoclakis S, Rossillo A, Lakkireddy D, Reddy M, Hao S, Hongo R, Beheiry S, Zagrodzky J, Rong B, Mohanty S, Elayi CS, Forleo G, Pelargonio G, Narducci ML, Dello Russo A, Casella M, Fassini G, Tondo C, Schweikert RA, Natale A. Periprocedural Stroke and Bleeding Complications in Patients Undergoing Catheter Ablation of Atrial Fibrillation With Different Anticoagulation Management: Results From the Role of Coumadin in Preventing Thromboembolism in Atrial Fibrillation (AF) Patients Undergoing Catheter Ablation (COMPARE) Randomized Trial. *Circulation* 2014;**129**(25):2638-44.
8. Cappato R, Marchlinski FE, Hohnloser SH, Naccarelli GV, Xiang J, Wilber DJ, Ma CS, Hess S, Wells DS, Juang G, Vijgen J, Hugl BJ, Balasubramaniam R, De Chillou C, Davies DW, Fields LE, Natale A, Investigators V-A. Uninterrupted rivaroxaban vs. uninterrupted vitamin K antagonists for catheter ablation in non-valvular atrial fibrillation. *Eur Heart J* 2015;**36**(28):1805-11.
9. Calkins H, Willems S, Gerstenfeld EP, Verma A, Schilling R, Hohnloser SH, Okumura K, Serota H, Nordaby M, Guiver K, Biss B, Brouwer MA, Grimaldi M, Investigators R-C. Uninterrupted Dabigatran versus Warfarin for Ablation in Atrial Fibrillation. *N Engl J Med* 2017;**376**(17):1627-1636.
10. Medi C, Evered L, Silbert B, The A, Halloran K, Morton J, Kistler P, Kalman J. Subtle Post-Procedural Cognitive Dysfunction following Atrial Fibrillation Ablation. *J Am Coll Cardiol* 2013;**62**(6):531-9.
11. Schwarz N, Kuniss M, Nedelmann M, Kaps M, Bachmann G, Neumann T, Pitschner HF, Gerriets T. Neuropsychological decline after catheter ablation of atrial fibrillation. *Heart Rhythm* 2010;**7**(12):1761-7.

12. Haeusler KG, Kirchhof P, Endres M. Left atrial catheter ablation and ischemic stroke. *Stroke* 2012;**43**(1):265-70.
13. Wu S, Yang YM, Zhu J, Wan HB, Wang J, Zhang H, Shao XH. Meta-Analysis of Efficacy and Safety of New Oral Anticoagulants Compared With Uninterrupted Vitamin K Antagonists in Patients Undergoing Catheter Ablation for Atrial Fibrillation. *Am J Cardiol* 2016;**117**(6):926-34.
14. Herm J, Fiebach JB, Koch L, Kopp UA, Kunze C, Wollboldt C, Brunecker P, Schultheiss HP, Schirdewan A, Endres M, Haeusler KG. Neuropsychological effects of MRI-detected brain lesions after left atrial catheter ablation for atrial fibrillation: long-term results of the MACPAF study. *Circ Arrhythm Electrophysiol* 2013;**6**(5):843-50.
15. Benameur K, Bykowski JL, Luby M, Warach S, Latour LL. Higher prevalence of cortical lesions observed in patients with acute stroke using high-resolution diffusion-weighted imaging. *AJNR Am J Neuroradiol* 2006;**27**(9):1987-9.
16. Di Biase L, Callans D, Haeusler KG, Hindricks G, Al-Khalidi H, Mont L, Cosedis Nielsen J, Piccini JP, Schotten U, Kirchhof P. Rationale and design of AXAFA-AFNET 5: an investigator-initiated, randomized, open, blinded outcome assessment, multi-centre trial to comparing continuous apixaban to vitamin K antagonists in patients undergoing atrial fibrillation catheter ablation. *Europace* 2017;**19**(1):132-138.
17. Goette A, Schon N, Kirchhof P, Breithardt G, Fetsch T, Hausler KG, Klein HU, Steinbeck G, Wegscheider K, Meinertz T. Angiotensin II-antagonist in paroxysmal atrial fibrillation (ANTIPAF) trial. *Circ Arrhythm Electrophysiol* 2012;**5**(1):43-51.
18. Kirchhof P, Andresen D, Bosch R, Borggrefe M, Meinertz T, Parade U, Ravens U, Samol A, Steinbeck G, Treszl A, Wegscheider K, Breithardt G. Short-term versus long-term antiarrhythmic drug treatment after cardioversion of atrial fibrillation (Flec-SL): a prospective, randomised, open-label, blinded endpoint assessment trial. *Lancet* 2012;**380**(9838):238-46.
19. Nasreddine ZS, Phillips N, Chertkow H. Normative data for the Montreal Cognitive Assessment (MoCA) in a population-based sample. *Neurology* 2012;**78**(10):765-6; author reply 766.
20. Granger CB, Alexander JH, McMurray JJ, Lopes RD, Hylek EM, Hanna M, Al-Khalidi HR, Ansell J, Atar D, Avezum A, Bahit MC, Diaz R, Easton JD, Ezekowitz JA, Flaker G, Garcia D, Geraldes M, Gersh BJ, Golitsyn S, Goto S, Hermosillo AG, Hohnloser SH, Horowitz J, Mohan P, Jansky P, Lewis BS, Lopez-Sendon JL, Pais P, Parkhomenko A, Verheugt FW, Zhu J, Wallentin L. Apixaban versus warfarin in patients with atrial fibrillation. *N Engl J Med* 2011;**365**(11):981-92.
21. Le Heuzey JY, Ammentorp B, Darius H, De Caterina R, Schilling RJ, Schmitt J, Zamorano JL, Kirchhof P. Differences among western European countries in anticoagulation management of atrial fibrillation. Data from the PREFER IN AF registry. *Thromb Haemost* 2014;**111**(5):833-41.
22. Rosendaal FR, Cannegieter SC, van der Meer FJ, Briet E. A method to determine the optimal intensity of oral anticoagulant therapy. *Thromb Haemost* 1993;**69**(3):236-9.
23. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, Kaul S, Wiviott SD, Menon V, Nikolsky E, Serebruany V, Valgimigli M, Vranckx P, Taggart D, Sabik JF, Cutlip DE, Krucoff MW, Ohman EM, Steg PG, White H. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the bleeding academic research consortium. *Circulation* 2011;**123**(23):2736-47.
24. Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of the S, Standardization Committee of the International Society on T, Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 2005;**3**(4):692-4.
25. Arbelo E, Brugada J, Hindricks G, Maggioni AP, Tavazzi L, Vardas P, Laroche C, Anselme F, Inama G, Jais P, Kalarus Z, Kautzner J, Lewalter T, Mairesse GH, Perez-Villacastin J, Riahi S, Taborsky M, Theodorakis G, Trines SA, Atrial Fibrillation Ablation Pilot Study I. The atrial fibrillation ablation pilot study: a European Survey on Methodology and results of catheter ablation for atrial fibrillation conducted by the European Heart Rhythm Association. *Eur Heart J* 2014;**35**(22):1466-78.
26. Di Biase L, Briceno DF, Trivedi C, Mohanty S, Gianni C, Burkhardt JD, Mohanty P, Bai R, Gunda S, Horton R, Bailey S, Sanchez JE, Al-Ahmad A, Hranitzky P, Gallingshouse GJ, Reddy YM, Zagrodzky J, Hongo R, Beheiry S, Lakkireddy D, Natale A. Is transesophageal echocardiogram

mandatory in patients undergoing ablation of atrial fibrillation with uninterrupted novel oral anticoagulants? Results from a prospective multicenter registry. *Heart Rhythm* 2016;**13**(6):1197-202.

27. Haines DE, Stewart MT, Barka ND, Kirchhof N, Lentz LR, Reinking NM, Urban JF, Halimi F, Deneke T, Kanal E. Microembolism and Catheter Ablation II: Effects of Cerebral Microemboli Injection in a Canine Model. *Circ Arrhythm Electrophysiol* 2013;**6**(1):23-30.

28. Takami M, Lehmann HI, Parker KD, Welker KM, Johnson SB, Packer DL. Effect of Left Atrial Ablation Process and Strategy on Microemboli Formation During Irrigated Radiofrequency Catheter Ablation in an In Vivo Model. *Circ Arrhythm Electrophysiol* 2016;**9**(1):e003226.

29. Kirchhof P, Breithardt G, Bax J, Benninger G, Blomstrom-Lundqvist C, Boriani G, Brandes A, Brown H, Brueckmann M, Calkins H, Calvert M, Christoffels V, Crijns H, Dobrev D, Ellinor P, Fabritz L, Fetsch T, Freedman SB, Gerth A, Goette A, Guasch E, Hack G, Haegeli L, Hatem S, Haeusler KG, Heidebuchel H, Heinrich-Nols J, Hidden-Lucet F, Hindricks G, Juul-Moller S, Kaab S, Kappenberger L, Kespohl S, Kotecha D, Lane DA, Leute A, Lewalter T, Meyer R, Mont L, Munzel F, Nabauer M, Nielsen JC, Oeff M, Oldgren J, Oto A, Piccini JP, Pilmeyer A, Potpara T, Ravens U, Reinecke H, Rostock T, Rustige J, Savelieva I, Schnabel R, Schotten U, Schwichtenberg L, Sinner MF, Steinbeck G, Stoll M, Tavazzi L, Themistoclakis S, Tse HF, Van Gelder IC, Vardas PE, Varpula T, Vincent A, Werring D, Willems S, Ziegler A, Lip GY, Camm AJ. A roadmap to improve the quality of atrial fibrillation management: proceedings from the fifth Atrial Fibrillation Network/European Heart Rhythm Association consensus conference. *Europace* 2016;**18**(1):37-50.

Legends to Figures

Figure 1: CONSORT diagram of the AXAFA – AFNET 5 study

Figure 2A: Cumulative primary outcome events since randomization until 90 days after randomization at full scale (upper panel) and magnified (lower panel) in the ablation set. VKA vitamin K antagonist therapy.

Figure 2B: Cumulative primary outcome events starting from ablation until 90 days after ablation at full scale (upper panel) and magnified (lower panel). VKA vitamin K antagonist therapy.

Figure 3: Forest plot of the differences (90% confidence intervals) in event rates in the main clinical subgroups. VKA vitamin K antagonist therapy.

Figure 4: Examples of acute brain lesions detected in the brain magnetic resonance imaging sub-study. Acute brain lesions (arrows in Figures 4A, 4C, 4E, 4G) are found by high resolution diffusion weighted brain magnetic resonance imaging (DWI) without corresponding lesions in the Fluid-attenuated inversion recovery (FLAIR) images (Figure 4B, 4D, 4F, 4H). Shown are representative lesions in two patients randomized to apixaban (Figure 4A/B; 4C/D) and in two patients randomized to VKA (Figure 4E/F; 4 G/H). The FLAIR images also detected chronic white matter lesions (asterisks (*) in Figure 4F).

Summary Figure. Cumulative outcome events in AXAFA – AFNET 5 in patients undergoing atrial fibrillation ablation at risk of stroke (top) and comparison to event rates in the two other controlled trials comparing continuous NOAC therapy with continuous VKA therapy (bottom). TIMI major bleeds were not separately reported in the main paper of RE-CIRCUIT, ISTH major bleeds were not separately reported in the main paper of VENTURE-AF.

Table 1: Inclusion and exclusion criteria of the AXAFA – AFNET 5 trial. Reproduced from the AXAFA – AFNET 5 design paper¹⁶.

Inclusion	Exclusion
Non-valvular atrial fibrillation (ECG-documented) with a clinical indication for catheter ablation	Any disease that limits life expectancy to less than 1 year
Clinical indication to undergo catheter ablation on continuous anticoagulant therapy	Participation in another clinical trial, either within the past 2 months or still ongoing
Presence of at least one of the CHADS ₂ stroke risk factors*	Previous participation in AXAFA
Age \geq 18 years	Pregnant women or women of childbearing potential not on adequate birth control: only women with a highly effective method of contraception (oral contraception or intra-uterine device) or sterile women can be randomized
Provision of signed informed consent	Breastfeeding women
	Drug abuse or clinically manifest alcohol abuse
	Any stroke within 14 days before randomization
	Concomitant treatment with drugs that are strong dual inhibitors of cytochrome P450 3A4 (CYP3A4) and P-glycoprotein (P-gp) or strong dual inducers of CYP3A4 and P-gp
	Valvular AF (as defined by the focused update of the ESC guidelines on AF, i.e. severe mitral valve stenosis, mechanical heart valve). Furthermore, patients who underwent mitral valve repair are not eligible for AXAFA
	Any previous ablation or surgical therapy for AF
	Cardiac ablation therapy for any indication (catheter-based or surgical) within 3 months prior to randomization
	Clinical need for “triple therapy” (combination therapy of clopidogrel, acetylsalicylic acid, and oral anticoagulation)
	Other contraindications for use of VKA or apixaban
	Documented atrial thrombi less than 3 months prior to randomization
	Severe chronic kidney disease with an estimated glomerular filtration rate (GFR) $<$ 15 ml/min

*Stroke or TIA, age \geq 75 years, hypertension, defined as chronic treatment for hypertension, estimated need for continuous antihypertensive therapy or resting blood pressure $>$ 145/90 mm Hg, diabetes mellitus, symptomatic heart failure (NYHA \geq II).

Table 2: Clinical characteristics of the AXAFA – AFNET 5 ablation population.

Number of patients with valid information (n (%)) is only given when values were missing.

BD twice daily dosing; SD standard deviation; q1, q3 are 25th and 75th percentiles, respectively; VKA vitamin K antagonist.

	All patients n=633	Apixaban n=318 (n=317 5 mg BD, n=1 2.5 mg BD)	VKA n=315 (n=127 warfarin, n=102 phenprocoumon, n=86 acenocoumarol)
Age (years)			
Median age (q1, q3)	64 (58, 70)	64 (57, 70)	64 (58, 70)
Female sex	209 (33%)	100 (31%)	109 (35%)
Weight (kg)			
Median weight (q1, q3)	87.0 (77.0, 99.3)	88.0 (77.0, 100.0)	86.6 (76.0, 98.0)
Median Body Mass Index (q1, q3)	28.3 (25.3, 31.6)	28.4 (25.5, 31.3)	28.2 (25.2, 31.9)
Concomitant conditions, stroke risk factors and CHA₂DS₂VASc score			
CHA ₂ DS ₂ VASc score, mean (SD)	2.4 (1.2)	2.4 (1.2)	2.4 (1.2)
CHA ₂ DS ₂ VASc score, median (q1, q3)	2 (2, 3)	2 (1, 3)	2 (2, 3)
Hypertension (n)	571 (90.2%)	283 (89.0%)	288 (91.4%)
Median systolic blood pressure (q1, q3)	138.0 (125.0, 150.0)	137.0 (125.0, 149.5)	140.0 (125.0, 152.0)
Median diastolic blood pressure (q1, q3)	82.0 (76.0, 90.0)	82.0 (75.0, 91.0)	82.0 (77.0, 90.0)
Symptomatic heart failure (NYHA II-IV)	150 (23.7%)	78 (24.5%)	72 (22.9%)
NYHA I	62 (9.8%)	30 (9.4%)	32 (10.2%)
NYHA II	126 (19.9%)	67 (21.1%)	59 (18.7%)
NYHA III	24 (3.8%)	11 (3.5%)	13 (4.1%)
NYHA IV	0	0	0
Diabetes mellitus	76 (12.0%)	41 (12.9%)	35 (11.1%)
Prior stroke or transient ischemic attack	47 (7.4%)	24 (7.5%)	23 (7.3%)
Age ≥ 75 years	56 (8.8%)	28 (8.8%)	28 (8.9%)
Age 65 – 74 years	240 (37.9%)	122 (38.4%)	118 (37.5%)
Vascular disease, defined as coronary artery disease, peripheral artery disease, or carotid disease	83 (13.1%)	41 (12.9%)	42 (13.3%)
Valvular heart disease	73 (11.5%)	39 (12.3%)	34 (10.8%)
Mitral valve disease (moderate or more)	20	15	5
Aortic valve disease (moderate or more)	6	3	3

	All patients	Apixaban	VKA
Confirmed coronary artery disease	77 (12.2%)	39 (12.3%)	38 (12.1%)
Chronic obstructive lung disease	39 (6.2%)	21 (6.6%)	18 (5.7%)
Clinical history of major bleeding	13 (2.1%)	10 (3.1%)	3 (1.0%)
Concomitant medical therapy			
n (%)	633	318	315
Amiodarone	102 (16.1%)	49 (15.4%)	53 (16.8%)
Dronedarone	13 (2.1%)	3 (0.9%)	10 (3.2%)
Flecainide	125 (19.7%)	59 (18.6%)	66 (21.0%)
Propafenone	16 (2.5%)	8 (2.5%)	8 (2.5%)
Sotalol >160mg/d	16 (2.5%)	7 (2.2%)	9 (2.9%)
ACE inhibitor or angiotensin receptor blocker	388 (61.3%)	192 (60.4%)	196 (62.2%)
Calcium channel antagonists	147 (23.2%)	72 (22.6%)	75 (23.8%)
Diuretics	221 (34.9%)	120 (37.7%)	101 (32.1%)
Antianginal medication	2 (0.3%)	0	2 (0.6%)
Antidiabetic medication	63 (10.0%)	32 (10.1%)	31 (9.8%)
Statins	231 (36.5%)	111 (34.9%)	120 (38.1%)
Platelet inhibitors or non-steroidal anti-inflammatory agents	30 (4.7%)	11 (3.5%)	19 (6.0%)
Beta blockers	451 (71.2%)	230 (72.3%)	221 (70.2%)
Digoxin or digitoxin	26 (4.1%)	17 (5.3%)	9 (2.9%)
Last INR before ablation (n)	531 (83.9)	217 (68.2%)	314 (99.7%)
Mean (SD)	1.9 (0.7)	1.2 (0.3), P<0.001 vs. VKA	2.3 (0.5)
Median (q1, q3)	2.0 (1.1, 2.4)	1.1 (1.0, 1.2)	2.3 (2.0, 2.6)
Quality of Life at baseline			
SF-12 physical component, n (%)	44.6 (37.7, 51.4), n=597	43.5 (38.1, 51.3), n=301	45.2 (37.6, 51.5), n=296
SF-12 mental component n (%)	598 (94.5%)	301 (94.7%)	297 (94.3%)
SF-12 mental component n (%)	50.3 (42.8, 57.5), n=598	51.2 (43.0, 57.9), n=301	49.7 (42.6, 57.4), n=297
Karnofsky scale	90 (80, 90)	80 (80, 90)	90 (80, 90)
Cognitive function (Montreal Cognitive Assessment, MoCA) at baseline			
Median MoCA(q1, q3)	27.0 (25.0, 29.0), n=618	27.0 (25.0, 29.0), n=313	27.0 (25.0, 29.0), n=305
At least mild cognitive impairment (MoCA <26)	188 (30.4%)	93 (29.7%)	95 (31.1%)
Modified EHRA scale at baseline			
mEHRA I	40 (6.3%)	18 (5.7%)	22 (7.0%)
mEHRA IIa	164 (25.9%)	76 (23.9%)	88 (27.9%)
mEHRA IIb	205 (32.4%)	107 (33.6%)	98 (31.1%)
mEHRA III	208 (32.9%)	110 (34.6%)	98 (31.1%)
mEHRA IV	16 (2.5%)	7 (2.2%)	9 (2.9%)
Ablation information			
Atrial fibrillation pattern			
Paroxysmal atrial fibrillation	367 (58.0%)	189 (59.4%)	178 (56.5%)

	All patients	Apixaban	VKA
Persistent or long-standing persistent atrial fibrillation	266 (42.0%)	129 (40.6%)	137 (43.5%)
Time from randomization to ablation (days)			
Mean (SD)	38.0 (27.3)	36.9 (27.6)	39.1 (27.0)
Median (q1, q3)	35.0 (20.0, 50.0)	34.0 (18.0, 48.0)	36.0 (21.0, 52.0)
Rhythm at start of ablation			
Number of patients	633	318	315
Sinus rhythm	434 (68.6%)	212 (66.6%)	222 (70.6%)
Atrial fibrillation	180 (28.4%)	98 (30.8%)	82 (26.0%)
Atrial flutter	12 (1.9%)	3 (0.9%)	9 (2.8%)
Pacing	7 (1.1%)	5 (1.6%)	2 (0.6%)
Other	0 (0%)	0 (0%)	0 (0%)
Type of ablation			
Pulmonary vein isolation, n (%)	571 (90.2%)	288 (90.6%)	283 (89.8%)
Pulmonary vein isolation plus other ablation, n (%)	59 (9.3%)	29 (9.1%)	30 (9.5%)
Other ablation without pulmonary vein isolation	3 (0.5%)	1 (0.3%)	2 (0.6%)
Transesophageal echocardiography prior to ablation	549 (86.7%)	269 (84.6%)	280 (88.9%)
Total duration of ablation procedure (minutes), Median (q1, q3)			
	135 (110, 175)	136 (110, 175)	135 (105, 172)
Ablation energy source			
Radio frequency	402 (63.5%)	207 (65.1%)	195 (61.9%)
Cryoablation	186 (29.3%)	92 (28.9%)	94 (29.8%)
Other	45 (7.1%)	19 (6.0%)	26 (8.3%)
Abnormal blood parameters			
Red blood cell count Abnormal	65 / 618 (10.5%)	32 / 311 (10.3%)	33 / 307 (10.7%)
Platelet count abnormal	35 / 625 (5.6%)	20 / 315 (6.3%)	15 / 310 (4.8%)
ALT abnormal	75 / 612 (12.3%)	39 / 307 (12.7%)	36 / 305 (11.8%)
Bilirubin abnormal	38 / 596 (6.4%)	14 / 297 (4.7%)	24 / 299 (8.0%)

Table 3: Primary outcomes in the AXAFA – AFNET 5 trial (ablation set), including details of the type of bleeding. Shown are number of patients per group. Some patients had more than one event. BARC4 events were not observed in the study. BD twice daily dosing.

	All patients	Apixaban	VKA
Patients with primary endpoint: composite of all-cause death, stroke or major bleeding	45/633 (7.1%)	22/318 (6.9%), non-inferiority p=0.0002	23/315 (7.3%)
Death	2 (0.3%)	1 (0.3%)	1 (0.3%)
Stroke or TIA	2 (0.3%)	2 (0.6%)	0
Major bleeding (BARC 2-5)	45 (7.1%)	20 (6.2%)	25 (7.9%)
<i>Bleeding requiring medical attention (BARC 2)</i>	<i>24 (3.8%)</i>	<i>12 (3.7%)</i>	<i>12 (3.8%)</i>
<i>Bleeding with hemoglobin drop of 30 to <50 g/l or requiring transfusion (BARC 3a)</i>	<i>9 (1.4%)</i>	<i>5 (1.6%)</i>	<i>4 (1.3%)</i>
<i>Bleeding with hemoglobin drop ≥50 g/l, or requiring surgery or iv vasoactive agents, or cardiac tamponade (BARC 3b)</i>	<i>11 (1.7%)</i>	<i>3 (0.9%)</i>	<i>8 (2.5%)</i>
<i>Intracranial hemorrhage (BARC 3c)</i>	<i>1 (0.2%)</i>	<i>0</i>	<i>1 (0.3%, fatal)</i>
TIMI major bleeding (Intracranial bleed, or bleeding resulting in a hemoglobin drop of ≥ 50 g/l, or bleeding resulting in death within 7 days)	4 (0.6%)	1 (0.3%)	3 (1%)
ISTH major bleeding	24 (3.8%)	10 (3.1%)	14 (4.4%)
Bleeding event by clinical type			
Tamponade	7 (1.1%)	2 (0.6%)	5 (1.6%)
Access site bleed	27 (4.3%)	12 (3.8%)	15 (4.8%)
Bleeding requiring transfusion of red blood cells	3 (0.5%)	2 (0.6%)	1 (0.3%)
Other major bleed	7 (1.1%)	5 (1.6%)	2 (0.6%)

Table 4: Secondary outcomes in the AXAFA – AFNET 5 trial (ablation set).

Number of patients with valid information (n (%)) is only given when values were missing. P values marked by asterisks (*) indicate differences between baseline and end of follow-up measurements. BD twice daily dosing; q1, q3 indicate 25th and 75th percentiles, respectively.

	All patients n=633	Apixaban n=318 (n=317 5 mg BD, n=1 2.5 mg BD)	VKA n=315 (n=127 warfarin, n=102 phenprocoumon, n=86 acenocoumarol)
time from randomization to ablation in days, median (q1, q3)	35.0 (20.0, 50.0)	34.0 (18.0, 48.0)	36.0 (21.0, 52.0)
Nights spent in hospital after index ablation, median (q1, q3)	3 (2, 5)	2 (1, 5)	3 (2, 7)
ACT during ablation in seconds, median (q1, q3)	325.0 (285.0, 370.0)	310.0 (273.0, 350.0)	348.5 (304.0, 396.0)
Number of subjects with all ACT values in range (n, %)	234 / 631 (37.1%)	73 / 316 (23.1%)	161 / 315 (51.1%)
Number of subjects with at least one ACT value < 250 (n, %)	214 / 631 (33.9%)	130 / 316 (41.1%)	84 / 315 (26.7%)
Number of subjects with at least one ACT value < 300 (n, %)	397 / 631 (62.9%)	243 / 316 (76.9%)	154 / 315 (48.9%)
Number of bleeding events (n)	118	54	64
Patients without recurrence of atrial fibrillation (n, %)	434 / 619 (70.1%)	217 / 311 (69.8%)	217 / 308 (70.5%)
Quality of life			
SF-12 physical component score at end of study, median (q1, q3), n	48.6 (42.0, 54.2), n=564	48.4 (41.9, 54.2), n=289	48.8 (42.2, 54.4), n=275
Change in SF-12 physical component score at end of study compared to baseline, median (q1, q3), n	2.5 (-2.1, 8.1), n=547, p<0.001*	2.4 (-2.2, 7.9), n=280	2.8 (-2.0, 8.3), n=267
SF-12 mental component score at end of study, median (q1, q3), n	54.4 (46.0, 58.6), n=565	54.2 (45.8, 58.3), n=290	54.5 (46.6, 59.7), n=267
Change in SF-12 mental component score at end of study compared to baseline, median (q1, q3), n	1.2 (-3.2, 8.0), n=548, p<0.001*	0.4 (-3.6, 8.0), n=281	1.6 (-2.8, 8.3), n=267
Karnofsky score at end of study, median (q1, q3), n	100 (90, 100), n=619	100 (90, 100), n=311	100 (90, 100), n=308
Change in Karnofsky score at end of study compared to	10 (0, 10), n=619	10 (0, 10), n=311	10 (0, 10), n=308

baseline (Δ Karnofsky), median (q1, q3),			
Cognitive Function (Montreal Cognitive Assessment, MoCA)			
Cognitive function at end of study (MoCA), median (q1, q3), n	28.0 (26.0, 29.0), n=607	28.0 (26.0, 29.0), n=305	28.0 (26.0, 29.0), n=302
Abnormal MoCA at baseline (<26), n (%)	141 (23.2%)	75 (24.6%)	66 (21.9%)
Change in MoCA at end of study compared to baseline, Median (q1, q3), n	1.0 (-1.0, 2.0), n=597, $p<0.001^*$	0.0 (-1.0, 2.0), n=301	1.0 (-1.0, 2.0), n=296
Change in patients with abnormal MoCA at end of study compared to baseline, n (%)	141/607 (23.2%), -7.2%, $p=0.005^*$	75/305 (24.6%) -5.1%	66/302 (21.9%) -9.2%

Table 5: Acute brain lesions detected by high resolution diffusion-weighted magnetic resonance imaging (MRI sub-study). P values were determined by Pearson's chi square test.

	All patients (n=323)	Apixaban (n=162)	VKA (n=161)	P value
No lesion	239 (74.0%)	118 (72.8%)	121 (75.2%)	0.635
Exactly one lesion	46 (14.2%)	27 (16.7%)	19 (11.8%)	0.211
Exactly two lesions	21 (6.5%)	7 (4.3%)	14 (8.7%)	0.111
More than two lesions	17 (5.3%)	10 (6.2%)	7 (4.3%)	0.463

Figure 1






Figure 2A top panel



AXAFA-AFNET 5
EudraCT no. 2014-002442-45


Figure 4b
Kaplan-Meier Cumulative Event Rate of Composite Primary Endpoint Magnified
(mITT Population¹)



¹mITT population: represents all randomized patients who underwent index catheter ablation

AXAFA-AFNET 5
EudraCT no. 2014-002442-45


Figure 4g
Kaplan-Meier Cumulative Event Rate of Composite Primary Endpoint
(mITT Population¹)



¹mITT population: represents all randomized patients who underwent index catheter ablation

AXAFA-AFNET 5
EudraCT no. 2014-002442-45


Figure 4h
Kaplan-Meier Cumulative Event Rate of Composite Primary Endpoint Magnified
(mITT Population¹)



¹mITT population: represents all randomized patients who underwent index catheter ablation





AXAFA-AFNET 5
EudraCT no. 2014-002442-45

Figure 5
Forest Plot of Primary Composite Event Rate Differences with Respect to Covariate Subgroups
(mITT Population¹)



¹mITT population: represents all randomized patients who underwent index catheter ablation
²87.0 kg is the median weight in the ITT population

Figure 4



Cumulative Outcome Events in AXAFA – AFNET 5 (top) and event rates in the patients undergoing ablation the three trials comparing continuous NOAC to VKA in AF ablation patients (bottom)

TIMI major bleeds were not reported in RE-CIRCUIT
ISTH major bleeds were not reported in VENTURE-AF

